


2 通院中治療者(医療機関[糖尿病外来]受診者) 結果

性別・年齢別人数

性別	年齢	集計	集計
男性	30歳代	2	10
	40歳代	3	
	50歳代	5	
	60歳代	19	
	70歳代	24	
男性計		53	53
女性	30歳代	2	8
	40歳代	1	
	50歳代	5	
	60歳代	10	
	70歳代	8	
女性計		26	26
総計		79	79



性別・年代別・罹患疾病(糖尿病)

性別	年齢	治療中糖尿病	服薬あり	服薬なし	総計	服薬あり	服薬なし
男性	30~50歳代	糖尿病	7	2	9	77.8%	22.2%
	60歳代	糖尿病	14	3	17	82.4%	17.6%
	70歳代	糖尿病	18	0	18	100.0%	0.0%
男性計			39	5	44	88.6%	11.4%
女性	30~50歳代	糖尿病	5	3	8	62.5%	37.5%
	60歳代	糖尿病	9	0	9	100.0%	0.0%
	70歳代	糖尿病	8	0	8	100.0%	0.0%
女性計			22	3	25	88.0%	12.0%
総計			62	8	70	88.6%	11.4%

性別・年代別・罹患疾病(高血圧)

性別	年齢	治療中糖尿病	治療中高血圧	服薬あり	総計	服薬あり
男性	30-50歳代	糖尿病	高血圧	3	3	100.0%
	60歳代	糖尿病	高血圧	5	5	100.0%
	70歳代	糖尿病	高血圧	4	4	100.0%
男性計				12	12	100.0%
女性	60歳代	糖尿病	高血圧	2	2	100.0%
	70歳代	糖尿病	高血圧	2	2	100.0%
女性計				4	4	100.0%
総計				16	16	100.0%

性別・年齢別・罹患疾病(糖尿病・高血圧以外)

性別	年齢	治療中糖尿病	その他内容治療中の疾病	服薬あり	総計	服薬あり
男性	30-50歳代	糖尿病	高脂血症	1	1	100.0%
			心筋梗塞	1	1	100.0%
			心臓病	1	1	100.0%
			不眠症	1	1	100.0%
		糖尿病計		4	4	100.0%
	60歳代	糖尿病	更年期	1	1	100.0%
			循環器	1	1	100.0%
			腎臓病	1	1	100.0%
		糖尿病計		3	3	100.0%
	70歳代	糖尿病	C型肝炎	1	1	100.0%
	糖尿病計		1	1	100.0%	
男性集計				8	8	100.0%
女性	30-50歳代	糖尿病	バセドウ病	1	1	100.0%
			糖尿病計	1	1	100.0%
女性集計				1	1	100.0%
総計				9	9	100.0%

就寝時間(主に寝る時間)*○は1つ

性別	年齢	治療中 糖尿病	その他内 容 治療中の	1.21時 前	2.21時 から22 時	3.22時 から23 時	4.23時 から24 時	5.24時 以降	総計	1.21時 前	2.21時 から22 時	3.22時 から23 時	4.23時 から24 時	5.24時 以降	
男性	30~50歳代	糖尿病	なし・空白	1	1	1	2	1	6	16.7%	16.7%	16.7%	33.3%	16.7%	
			高血圧症			1			1			100.0%			
			心筋梗塞						1	1					100.0%
			心臓病				1			1			100.0%		
			不眠症	1						1	100.0%				
	糖尿病計			2	1	3	2	2	10	20.0%	10.0%	30.0%	20.0%	20.0%	
	60歳代	糖尿病	なし・空白	1	3	5	4	1	14	7.1%	21.4%	35.7%	28.6%	7.1%	
			更年期		1				1		100.0%				
			循環器			1			1			100.0%			
			腎臓病		1				1		100.0%				
			糖尿病計		1	5	6	4	1	17	5.9%	29.4%	35.3%	23.5%	5.9%
	なし なし集計	なし・空白						1	1					100.0%	
								1	1					100.0%	
	70歳代	糖尿病	なし・空白	4	6	6	1		17	23.5%	35.3%	35.3%	5.9%		
			C型肝炎	1					1	100.0%					
糖尿病計			5	6	6	1		18	27.8%	33.3%	33.3%	5.6%			
男性集計			8	12	15	7	4	46	17.4%	26.1%	32.6%	15.2%	8.7%		
女性	30~50歳代	糖尿病	なし・空白			3		3	6			50.0%		50.0%	
			(糖尿病I型)					1	1					100.0%	
			バセドウ病		1				1		100.0%				
			糖尿病計		1	3		4	8		12.5%	37.5%		50.0%	
	60歳代	糖尿病	なし・空白	2	2	3	1	1	9	22.2%	22.2%	33.3%	11.1%	11.1%	
			糖尿病計	2	2	3	1	1	9	22.2%	22.2%	33.3%	11.1%	11.1%	
	70歳代	糖尿病	なし・空白		5	2	1		8		62.5%	25.0%	12.5%		
			糖尿病計		5	2	1		8		62.5%	25.0%	12.5%		
	女性集計			2	8	8	2	5	25	8.0%	32.0%	32.0%	8.0%	20.0%	
	総計				10	20	23	9	9	71	14.1%	28.2%	32.4%	12.7%	12.7%

就寝前の2時間以内に食べたり「カロリーのある飲みもの」を飲んだりするか*○は1つ

性別	年齢	治療中 糖尿病	その他内 容 治療中の	毎日	1日お き	1週間に2~3 回	1週間に1回 以下	未記入	総計	毎日	1日お き	1週間に2~3 回	1週間に1回 以下
男性	30~50歳代	糖尿病	なし・空白			5	1		6	83.3%	16.7%		
			高脂血症				1		1		100.0%		
			心筋梗塞			1			1		100.0%		
			心臓病			1			1		100.0%		
			不眠症		1				1	100.0%			
	糖尿病計			1	7	2		10	10.0%	70.0%	20.0%		
	60歳代	糖尿病	なし・空白	4	2	2	5	1	14	14.3%	14.3%	35.7%	7.1%
			更年期					1	1			100.0%	
			循環器					1	1			100.0%	
			腎臓病					1	1			100.0%	
			糖尿病計		4	2	2	8	1	17	11.8%	11.8%	47.1%
	なし なし集計	なし・空白				1		1		100.0%			
						1		1		100.0%			
	70歳代	糖尿病	なし・空白	2	2	4	8	1	17	11.8%	23.5%	47.1%	5.9%
			C型肝炎					1	1			100.0%	
男性集計				2	2	4	9	1	18	11.1%	22.2%	50.0%	5.6%
女性				6	5	14	19	2	46	10.9%	30.4%	41.3%	4.3%
女性	30~50歳代	糖尿病	なし・空白		1	1	4		6	16.7%	16.7%	66.7%	
			(糖尿病I型)				1		1		100.0%		
			バセドウ病					1	1			100.0%	
			糖尿病計		1	2	5		8	12.5%	25.0%	62.5%	
	60歳代	糖尿病	なし・空白	1		1	7		9		11.1%	77.8%	
			糖尿病計	1		1	7		9		11.1%	77.8%	
	70歳代	糖尿病	なし・空白	2		2	4		8		25.0%	50.0%	
			糖尿病計	2		2	4		8		25.0%	50.0%	
	女性集計				3	1	5	16		25	4.0%	20.0%	64.0%
	総計				9	6	19	35	2	71	8.5%	26.8%	49.3%

お通じ(排便)の頻度 *○は1つ


性別	年齢	治療中 糖尿病	その他内 容 治療中の	1.毎日	2.1日お き	3.1週間 に2~3 回	4.1週間 に1回 以下	未記入	総計	1.毎日	2.1日お き	3.1週間 に2~3 回	4.1週間 に1回 以下
男性	30~50歳代	糖尿病	なし・空白	4	1	1			6	66.7%	16.7%	16.7%	
			高脂血症			1			1		100.0%		
			心筋梗塞	1					1	100.0%			
			心臓病	1					1	100.0%			
			不眠症				1			1		100.0%	
		糖尿病計		6	2	2			10	60.0%	20.0%	20.0%	
	60歳代	糖尿病	なし・空白	8	2	4			14	57.1%	14.3%	28.6%	
			更年期			1			1		100.0%		
			循環器	1					1	100.0%			
			腎臓病	1					1	100.0%			
			糖尿病計		10	3	4			17	58.8%	17.6%	23.5%
		なし なし集計		1				1	100.0%				
	70歳代	糖尿病	なし・空白	9	6	1		1	17	52.9%	35.3%	5.9%	
			C型肝炎	1					1	100.0%			
			糖尿病計		10	6	1		1	18	55.6%	33.3%	5.6%
男性集計			27	11	7		1	46	58.7%	23.9%	15.2%		
女性	30~50歳代	糖尿病	なし・空白	1	2	2	1		6	16.7%	33.3%	33.3%	16.7%
			(糖尿病I型)			1			1		100.0%		
			バセドウ病	1					1	100.0%			
		糖尿病計		2	3	2	1		8	25.0%	37.5%	25.0%	12.5%
	60歳代	糖尿病	なし・空白	5	2	2			9	55.6%	22.2%	22.2%	
			糖尿病計		5	2	2		9	55.6%	22.2%	22.2%	
	70歳代	糖尿病	なし・空白	4	3	1			8	50.0%	37.5%	12.5%	
			糖尿病計		4	3	1		8	50.0%	37.5%	12.5%	
	女性集計			11	8	5	1		25	44.0%	32.0%	20.0%	4.0%
	総計				38	19	12	1	1	71	53.5%	26.8%	16.9%

健康を意識しているか *○は1つ

性別	年齢	治療中 糖尿病	その他内 容 治療中の	1.意識 してい る	2.意識 してい ない	未記入	総計	1.意識 してい る	2.意識 してい ない	未記入
男性	30~50歳代	糖尿病	なし・空白	3	2	1	6	50.0%	33.3%	16.7%
			高脂血症			1	1		100.0%	
			心筋梗塞		1		1		100.0%	
			心臓病	1			1	100.0%		
			不眠症	1			1	100.0%		
		糖尿病計		5	3	2	10	50.0%	30.0%	20.0%
	60歳代	糖尿病	なし・空白	10	3	1	14	71.4%	21.4%	7.1%
			更年期	1			1	100.0%		
			循環器	1			1	100.0%		
			腎臓病	1			1	100.0%		
			糖尿病計		13	3	1	17	76.5%	17.6%
		なし なし集計		1			1	100.0%		
	70歳代	糖尿病	なし・空白	13	4		17	76.5%	23.5%	
			C型肝炎	1			1	100.0%		
			糖尿病計		14	4		18	77.8%	22.2%
男性計			33	10	3	46	71.7%	21.7%	6.5%	
女性	30~50歳代	糖尿病	なし・空白	6			6	100.0%		
			(糖尿病I型)			1	1		100.0%	
			バセドウ病	1			1	100.0%		
		糖尿病計		7	1		8	87.5%	12.5%	
	60歳代	糖尿病	なし・空白	8	1		9	88.9%	11.1%	
			糖尿病計		8	1		9	88.9%	11.1%
	70歳代	糖尿病	なし・空白	7	1		8	87.5%	12.5%	
糖尿病計				7	1		8	87.5%	12.5%	
女性計			22	3		25	88.0%	12.0%		
総計			55	13	3	71	77.5%	18.3%	4.2%	

性別・年齢別 疾病名別 体格

性別	年齢	治療中 糖尿病	その他内容 治療中の疾 病	BMI20 未満	BMI20- 25未満	BMI25- 30未満	BMI30 以上	未記入	総計	BMI20 未満	BMI20- 25未満	BMI25- 30未満	BMI30 以上	
男性	30~50歳代	糖尿病	なし・空白			1	3	2	6			16.7%	50.0%	
			高脂血症			1			1			100.0%		
			心筋梗塞					1		1			100.0%	
			心臓病		1					1		100.0%		
			不眠症			1				1			100.0%	
		糖尿病計			1	3	4	2	10		10.0%	30.0%	40.0%	
	60歳代	糖尿病	なし・空白	1	4	7		2	14	7.1%	28.6%	50.0%		
			更年期			1			1			100.0%		
			循環器		1				1			100.0%		
			腎臓病			1			1			100.0%		
			糖尿病計		1	5	9	2	17	5.9%	29.4%	52.9%		
		なし			1			1			100.0%			
		なし集計			1			1			100.0%			
	70歳代	糖尿病	なし・空白	4	8	4		1	17	23.5%	47.1%	23.5%		
			C型肝炎					1	1					
糖尿病計			4	8	4	2	18	22.2%	44.4%	22.2%				
男性集計			5	15	16	4	6	46	10.9%	32.6%	34.8%	8.7%		
女性	30~50歳代	糖尿病	なし・空白		3	2	1	6		50.0%	33.3%	16.7%		
			(糖尿病I型)	1				1	100.0%					
			バセドウ病		1			1	100.0%					
			糖尿病計	1	4	2	1	8	12.5%	50.0%	25.0%	12.5%		
	60歳代	糖尿病	なし・空白	1	6	2		9	11.1%	66.7%	22.2%			
			糖尿病計	1	6	2		9	11.1%	66.7%	22.2%			
	70歳代	糖尿病	なし・空白		5	2		1	8		62.5%	25.0%		
			糖尿病計		5	2	1	8		62.5%	25.0%			
	女性集計			2	15	6	1	1	25	8.0%	60.0%	24.0%	4.0%	
	総計				7	30	22	5	7	71	9.9%	42.3%	31.0%	7.0%




性別・年齢別 疾病名別 栄養素摂取量推計(食塩)

性別	年齢	治療中 糖尿病	その他内容 治療中の疾 病	6-7g 未満	7-9g 未満	9-12g 未満	12-15 g未満	15-20 g未満	20g以 上	未集 計	総計	
男性	30~50歳代	糖尿病	なし・空白			1	1	2		2	6	
			高脂血症				1				1	
			心筋梗塞				1				1	
			心臓病							1	1	
			不眠症				1				1	
	糖尿病計					1	4	2	1	2	10	
	60歳代	糖尿病	なし・空白			1	5	5	1		2	14
			更年期				1					1
			循環器				1					1
			腎臓病				1					1
	糖尿病計				1	8	5	1		2	17	
	なし	なし集計	なし・空白						1			1
									1			1
	70歳代	糖尿病	なし・空白		1		3	3	7	2	1	17
			C型肝炎									1
糖尿病計			1		3	3	7	2	2	2	18	
男性計				1		5	15	15	4	6	46	
女性	30~50歳代	糖尿病	なし・空白		1	4	1				6	
			(糖尿病I型)			1					1	
			バセドウ病			1					1	
	糖尿病計			1	6	1					8	
	60歳代	糖尿病	なし・空白			5	2	2				9
			糖尿病計			5	2	2				9
	70歳代	糖尿病	なし・空白		3	1	1	2			1	8
			糖尿病計		3	1	1	2			1	8
女性計				4	12	4	4		1	25		
総計				1	4	17	19	19	4	7	71	

性別・年齢別 疾病名別 栄養素摂取量推計(食塩)

■ 6-7g未満 □ 7-9g未満 ■ 9-12g未満 ■ 12-15g未満 ■ 15-20g未満 ■ 20g以上




性別・年齢別 疾病名別 栄養素摂取量推計(炭水化物%エネルギー) :50-60%が目安

性別	年齢	治療中 糖尿病	その他内 容 治療中の 疾病	50% 未満	50- 60% 未満	60- 80% 未満	不明	総計	50% 未満	50- 60% 未満	60- 80% 未満	
男性	30~50歳代	糖尿病	なし・空白	1	3		2	6	16.7%	50.0%		
			高脂血症			1		1			100.0%	
			心筋梗塞			1		1			100.0%	
			心臓病			1		1			100.0%	
			不眠症	1				1	100.0%			
		糖尿病計			2	5	1	2	10	20.0%	50.0%	10.0%
	60歳代	糖尿病	なし・空白	2	3	7	2	14	14.3%	21.4%	50.0%	
			更年期	1				1	100.0%			
			循環器			1		1			100.0%	
			腎臓病			1		1			100.0%	
			糖尿病計		3	5	7	2	17	17.6%	29.4%	41.2%
		なし なし集計				1		1		100.0%		
	70歳代	糖尿病	なし・空白	6	5	5	1	17	35.3%	29.4%	29.4%	
			C型肝炎				1	1				
	糖尿病計		6	5	5	2	18	33.3%	27.8%	27.8%		
男性計				11	16	13	6	46	23.9%	34.8%	28.3%	
女性	30~50歳代	糖尿病	なし・空白 (糖尿病I型)	1	1	4		6	16.7%	16.7%	66.7%	
			バセドウ病			1		1		100.0%		
			糖尿病計		1	3	4		8	12.5%	37.5%	50.0%
	60歳代	糖尿病	なし・空白	3	3	3		9	33.3%	33.3%	33.3%	
			糖尿病計	3	3	3		9	33.3%	33.3%	33.3%	
	70歳代	糖尿病	なし・空白	2	3	2	1	8	25.0%	37.5%	25.0%	
			糖尿病計	2	3	2	1	8	25.0%	37.5%	25.0%	
	女性計			6	9	9	1	25	24.0%	36.0%	36.0%	
	総計				17	25	22	7	71	23.9%	35.2%	31.0%

性別・年齢別 疾病名別 栄養素摂取量推計(炭水化物%エネルギー)

■ 50%未満 □ 50-60%未満 ■ 60-80%未満




性別・年齢別 疾病名 栄養素摂取量推計(脂質%エネルギー) :20-25%が目安

性別	年齢	治療中 糖尿病	その他内容 治療中の疾 病	20%未満	20-25%未 満	25-30%未 満	30-40% 未満	40%以上	不明	総計		
男性	30~50歳代	糖尿病	なし・空白		2	2				2	6	
			高脂血症	1							1	
			心筋梗塞					1				1
			心臓病		1							1
			不眠症		1							1
		糖尿病計			1	4	2	1		2	10	
	60歳代	糖尿病	なし・空白	3	8			1		2	14	
			更年期		1							1
			循環器		1							1
			腎臓病					1				1
			糖尿病計		3	10		2		2		17
		なし なし集計						1			1	
	70歳代	糖尿病	なし・空白	5	5	2	4			1	17	
			C型肝炎							1		1
			糖尿病計		5	5	2	4		2		18
	男性計			9	19	4	8		6	46		
	女性	30~50歳代	糖尿病	なし・空白	1	2	1	1	1			6
				(糖尿病I型)		1						
バセドウ病						1						1
糖尿病計					1	3	2	1	1			8
60歳代		糖尿病	なし・空白	2	1	2	4				9	
			糖尿病計		2	1	2	4				9
			なし集計									
70歳代		糖尿病	なし・空白		2	3	2			1	8	
			糖尿病計			2	3	2		1		8
			なし集計									
女性計			3	6	7	7	1	1	25			
総計			12	25	11	15	1	7	71			

性別・年齢別 疾病名別 栄養素摂取量推計(脂質%エネルギー)

■ 20%未満 □ 20-25%未満 ■ 25-30%未満 ■ 30-40%未満 ■ 40%以上



信号の色(摂取基準充足率)別人数割合
基準: 摂取不足の回避を目的とする「推定平均必要量」(EAR)
半数以上が必要量を満たす量「推奨量」(RDA)

	性別	年齢	1	2	3	4	5	判定不可	総計	1	2	3
			赤信号 EARの1倍未満	黄信号 EARの1倍以上 & RDAの	青信号 RDAの0.95倍以上					赤信号 EARの1倍未満	黄信号 EARの1倍以上 & RDAの	青信号 RDAの0.95倍以上
カルシウム	男性	30~50歳代	3		5			2	10	37.5%		62.5%
		60歳代	10	1	6			2	19	58.8%	5.9%	35.3%
		70歳代	9	1	12			2	24	40.9%	4.5%	54.5%
		男性 集計		22	2	23		6	53	46.8%	4.3%	48.9%
	女性	30~50歳代	3	3	2				8	37.5%	37.5%	25.0%
		60歳代	4	2	4				10	40.0%	20.0%	40.0%
		70歳代	2	1	4			1	8	28.6%	14.3%	57.1%
		女性 集計		9	6	10		1	26	36.0%	24.0%	40.0%
	総計		31	8	33		7	79	43.1%	11.1%	45.8%	
鉄	男性	30~50歳代		1	7			2	10		12.5%	87.5%
		60歳代	1		16			2	19	5.9%		94.1%
		70歳代		1	21			2	24		4.5%	95.5%
		男性 集計		1	2	44		6	53	2.1%	4.3%	93.6%
	女性	30~50歳代	1	4	3				8	12.5%	50.0%	37.5%
		60歳代			10				10			100.0%
		70歳代			7			1	8			100.0%
		女性 集計		1	4	20		1	26	4.0%	16.0%	80.0%
	総計		2	6	64		7	79	2.8%	8.3%	88.9%	
ビタミンC	男性	30~50歳代	2	2	4			2	10	25.0%	25.0%	50.0%
		60歳代	5	2	10			2	19	29.4%	11.8%	58.8%
		70歳代	5		17			2	24	22.7%		77.3%
		男性 集計		12	4	31		6	53	25.5%	8.5%	66.0%
	女性	30~50歳代	2		6				8	25.0%		75.0%
		60歳代		2	8				10		20.0%	80.0%
		70歳代	1		6			1	8	14.3%		85.7%
		女性 集計		3	2	20		1	26	12.0%	8.0%	80.0%
	総計		15	6	51		7	79	20.8%	8.3%	70.8%	
葉酸	男性	30~50歳代	1	1	6			2	10	12.5%	12.5%	75.0%
		60歳代	1	1	15			2	19	5.9%	5.9%	88.2%
		70歳代			22			2	24			100.0%
		男性 集計		2	2	43		6	53	4.3%	4.3%	91.5%
	女性	30~50歳代	1		7				8	12.5%		87.5%
		60歳代			10				10			100.0%
		70歳代			7			1	8			100.0%
		女性 集計		1		24		1	26	4.0%		96.0%
	総計		3	2	67		7	79	4.2%	2.8%	93.1%	

基準: 「生活習慣病の予防のために現在の日本人が当面の目標とすべき摂取量」(DG)

	性別	年齢	1	2	3	4	5	判定不可	総計	1	2	3	4	5
			赤信号 EARの1倍 未満	黄信号 EARの1倍 以上& RDAの	青信号 RDAの 0.95倍以 上	黄信号 II DG上限の 1倍以上& DG上限の 0.95倍以 上	赤信号 II DG上限の 1.3倍以上			赤信号	黄信号	青信号	黄信号 II	赤信号 II
食物繊維	男性	30~50歳代	4	1	3			2	10	50.0%	12.5%	37.5%		
		60歳代	5	9	3			2	19	29.4%	52.9%	17.6%		
		70歳代	5	11	6			2	24	22.7%	50.0%	27.3%		
	男性 集計		14	21	12			6	53	29.8%	44.7%	25.5%		
	女性	30~50歳代	2	5	1				8	25.0%	62.5%	12.5%		
		60歳代	2	3	5				10	20.0%	30.0%	50.0%		
		70歳代	2	3	2			1	8	28.6%	42.9%	28.6%		
	女性 集計		6	11	8			1	26	24.0%	44.0%	32.0%		
総計			20	32	20		7	79	27.8%	44.4%	27.8%			
カリウム	男性	30~50歳代	2	2	4			2	10	25.0%	25.0%	50.0%		
		60歳代	1	10	6			2	19	5.9%	58.8%	35.3%		
		70歳代	3	4	15			2	24	13.6%	18.2%	68.2%		
	男性 集計		6	16	25			6	53	12.8%	34.0%	53.2%		
	女性	30~50歳代	1	3	4				8	12.5%	37.5%	50.0%		
		60歳代		5	5				10		50.0%	50.0%		
		70歳代	1	1	5			1	8	14.3%	14.3%	71.4%		
	女性 集計		2	9	14			1	26	8.0%	36.0%	56.0%		
総計			8	25	39		7	79	11.1%	34.7%	54.2%			
食塩 (ナトリウム)	男性	30~50歳代				1	7	2	10				12.5%	87.5%
		60歳代				1	16	2	19				5.9%	94.1%
		70歳代			1	4	17	2	24			4.5%	18.2%	77.3%
	男性 集計				1	6	40	6	53			2.1%	12.8%	85.1%
	女性	30~50歳代			1	1	6		8			12.5%	12.5%	75.0%
		60歳代					10		10					100.0%
		70歳代			1	3	3	1	8			14.3%	42.9%	42.9%
	女性 集計				2	4	19	1	26			8.0%	16.0%	76.0%
総計				3	10	59	7	79			4.2%	13.9%	81.9%	
総脂質	男性	30~50歳代		1	5	2		2	10		12.5%	62.5%	25.0%	
		60歳代	1	2	10	4		2	19	5.9%	11.8%	58.8%	23.5%	
		70歳代	2	8	7	5		2	24	9.1%	36.4%	31.8%	22.7%	
	男性 集計		3	11	22	11		6	53	6.4%	23.4%	46.8%	23.4%	
	女性	30~50歳代		1	4	1	2		8		12.5%	50.0%	12.5%	25.0%
		60歳代		2	2	6			10		20.0%	20.0%	60.0%	
		70歳代			2	5		1	8			28.6%	71.4%	
	女性 集計			3	8	12	2	1	26		12.0%	32.0%	48.0%	8.0%
総計			3	14	30	23	2	7	79	4.2%	19.4%	41.7%	31.9%	2.8%
飽和脂肪酸	男性	30~50歳代		2	3	2	1	2	10		25.0%	37.5%	25.0%	12.5%
		60歳代	1	2	13	1		2	19	5.9%	11.8%	76.5%	5.9%	
		70歳代	2	5	10	4	1	2	24	9.1%	22.7%	45.5%	18.2%	4.5%
	男性 集計		3	9	26	7	2	6	53	6.4%	19.1%	55.3%	14.9%	4.3%
	女性	30~50歳代			6	1	1		8			75.0%	12.5%	12.5%
		60歳代	1		5	4			10	10.0%		50.0%	40.0%	
		70歳代			3	4		1	8			42.9%	57.1%	
	女性 集計		1		14	9	1	1	26	4.0%		56.0%	36.0%	4.0%
総計			4	9	40	16	3	7	79	5.6%	12.5%	55.6%	22.2%	4.2%